



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكرو فيلم

بسم الله الرحمن الرحيم



HANAA ALY



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكروفيلم



شبكة المعلومات الجامعية التوثيق الإلكتروني والميكروفيلم



HANAA ALY



شبكة المعلومات الجامعية
التوثيق الإلكتروني والميكروفيلم

جامعة عين شمس التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغييرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار



HANAA ALY

**Evaluation of Role of Magnetic
Resonance Imaging in Detection and
Characterization of Soft Tissue masses
of the Hand and Wrist**

Thesis

*Submitted for partial fulfillment of
Master Degree in Radiodiagnosis*

By

Hythem Hashim Said Ahmed Ali

M.B.B.C.H

Faculty of Medicine Al Jazirah University.

Supervised by

Dr. Sahar Mohamed El Gaafary

Professor of Radiodiagnosis

Faculty of Medicine

Ain Shams University

Dr. Susan Adil Ali Abdul Rahim

Assistant Professor of Radiodiagnosis

Faculty of Medicine

Ain Shams University

*Faculty of Medicine
Ain Shams University*

2020



وَقُلْ رَبِّ
أَنْزِلْنِي مُنْزَلًا مُّبَارَكًا
وَأَنْتَ خَيْرُ الْمُنْزِلِينَ



Acknowledgment



*First of all, I would like to thank **ALLAH** who granted me the strength to accomplish this work.*

I have been honored and privileged to work under the supervision of such distinguished and eminent professors. It has truly been a chance of lifetime.

*I would like to express my deepest and ultimate gratitude to **Prof. Dr. Sahar Mohamed El Gaafary**, Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, who helped me to take the first step. His kind supervision and kind encouragement can't be denied.*

*My deepest gratitude goes to **Ass. Prof. Dr. Susan Adil Ali AbdulRahim**, Assistant Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University, for her patience and support. Her faithful encouragement, continuous guidance, and valuable advices that gave me the motive to utilize available resources in order to put this thesis in acceptable form.*

I would also like to thank my parents, my beautiful wife and my three beautiful kids, for their support and patience.

I'm also thankful to all my colleagues and my big family for their kind help and support. I'm also very grateful for all the patients who helped me to bring this thesis out.

I would like to express my hearty thanks to Egypt my second country to give me this chance. I will not forget.

Hythem Hashim Said Ahmed Ali

List of Contents

Title	Page No.
List of Tables	i
List of Figures	ii
Introduction	1
Aim of the Work	3
Review of Literature	
Wrist Anatomy	4
Hand Anatomy	13
MR Imaging Technique	25
MRI of Common Soft Tissue Masses of the Hand and Wrist	42
Patients and Methods	69
Results	73
Case Presentation	89
Discussion	119
Summary and Conclusion	130
References	132

List of Tables

Table No.	Title	Page No.
Table (1):	Classification of swellings arising in the hand and wrist.....	42
Table (2):	Imaging parameters of the used protocol of MR imaging of the wrist and hand.....	71
Table (3):	The number and frequency of the lesions and mean age of the patients in the study population.....	78
Table (4):	The frequency of different lesions according to MRI signal	80
Table (5):	Classification of lesions according to their location	81

List of Figures

Fig. No.	Title	Page No.
Fig. (1):	Carpal anatomy, three-dimensional (3D) CT images show the normal wrist	5
Fig. (2):	Distal radioulnar, radiocarpal, and mid carpal joints	6
Fig. (3):	The articular surface of the distal end of the radius and the adjacent triangular cartilage are exposed by removal of the carpal bones.....	6
Fig. (4):	Extensor Tendons with the Six Tendon Sheath Compartments.....	8
Fig. (5):	Wrist flexors	9
Fig. (6):	Anatomy of the carpal tunnel	10
Fig. (7):	Normal MR anatomy of the ulnar nerve.....	12
Fig. (8):	Midcarpal joint.....	13
Fig. (9):	Anatomy of the pulley system of the finger	18
Fig. (10):	Sagittal T1-weighted MR image shows the insertion of the central slip on the base of the middle phalanx and the insertion of the terminal tendon on the base of the distal phalanx	20
Fig. (11):	Transaxial T1-weighted spin-echo MR image (TR/TE, 500/14) shows normal findings of intrinsic hand muscles	22
Fig. (12):	Superficial structures of the left palm and wrist.....	24
Fig. (13):	A , Digitally subtracted hand arteriogram, dorsopalmar projection. B , Venous phase of hand arteriogram	24
Fig. (14):	The 5-inch general purpose receive-only surface coil	26
Fig. (15):	An eight-channel receive-only wrist array coil.....	27
Fig. (16):	Philips microscopy coils	28

List of Figures *(Cont...)*

Fig. No.	Title	Page No.
Fig. (17):	The microscopy coils used on the Philips units can be combined coil with other coils	29
Fig. (18):	T1 WIS coronal and axial showing planning of axial, sagittal and coronal images respectively.....	30
Fig. (19):	Axial localizer slice shows proper orientation of wrist.....	32
Fig. (20):	Coronal T1 WIS showing the hypointense bands of the flexor digitorum tendons.....	34
Fig. (21):	Normal anatomy	34
Fig. (22):	Coronal 3D GRE image shows the extensor capi ulnaris tendon	35
Fig. (23):	Lister's tubercle (dorsal tubercle) of the distal radius on a coronal PD FSE image	35
Fig. (24):	Sagittal T1with and without fat suppression showing the normal carpal bone	36
Fig. (25):	Sagittal T1 Fat suppression arthrographic images showing the normal flexor and extensor ligaments	37
Fig. (26):	Normal anatomy of the tendons	39
Fig. (27):	Axial T1 WIS showing the median nerve, FP (flexor pollicis longus) & AP (adductor pollicis) tendons.....	40
Fig. (28):	Axial T1 WIS showing the Flexor retinaculum, median nerve, and flexor digitorum rofundus & superficialis tendons	40
Fig. (29):	Axial T1 WIS showing some of the extensor compartments	41

List of Figures (Cont...)

Fig. No.	Title	Page No.
Fig. (30):	Axial T1 WIS showing carpal tunnel, flexor tendons, and median nerve.....	41
Fig. (31):	Ganglion cyst.....	44
Fig. (32):	(a) Dorsal ganglion. (b) Axial T1 weighted image demonstrating an intermediate signal mass.....	44
Fig. (33):	(GCT) of the tendon sheath, MR imaging: lobulated mass next to the proximal interphalangeal joint, relatively hypointense on T1W (a), with areas of T2W hypointensity (b) (hemosiderin deposition), with intense enhancement on post contrast.....	46
Fig. (34):	MRI (T2 fat sat) demonstrates low signal intensity fibrous band in the superficial subcutaneous tissue in the ring finger, which running parallel the flexor tendon.....	48
Fig. (35):	Lipoma and Coronal short tau inversion recovery.....	50
Fig. (36):	Fibrolipomatous hamartoma.....	51
Fig. (37):	Haemangioma.....	52
Fig. (38):	Microcystic lymphatic malformation.....	53
Fig. (39):	AVM.....	54
Fig. (40):	Schwannoma.....	56
Fig. (41):	Neurofibroma.....	57
Fig. (42):	Nail-bed glomus tumour.....	58
Fig. (43):	Periosteal chondroma.....	60
Fig. (44):	Synovial sarcoma in the dorsoradial aspect of the hand.....	61
Fig. (45):	Melanoma mets.....	62
Fig. (46):	Synovial hyperplasia.....	63

List of Figures *(Cont...)*

Fig. No.	Title	Page No.
Fig. (47):	Accessory abductor digiti minimi. Coronal T1 weighted image demonstrating an accessory muscle belly.....	65
Fig. (48):	Accessory palmaris longus muscle. Sagittal T1weighted image demonstrating an accessory muscle belly at the level of the wrist	65
Fig. (49):	Pulp space infection (felon) with underlying osteomyelitis.....	66
Fig. (50):	Lesion Characterization on the Basis of MR Images	68
Fig. (51):	The percentage of patients according to age.	74
Fig. (52):	The frequency and percentage of patients according to sex.....	75
Fig. (53):	The frequency and percentage of affection of the right, left hand and wrist joint.	76
Fig. (54):	The frequency and percentage of different lesions in the study population.....	77
Fig. (55):	The frequency of different lesions in relation to size.	78
Fig. (56):	The frequency of different lesions in related to the bones.	79
Fig. (57):	Axial T1 (A), axial T2 (B), axial DWI (C) Sagittal T2(D), of the right hand showing intermediate signal in, high signal	90
Fig. (58):	Coronal T1 (A), Coronal T2 (B), Coronal STIR (C) showing intermediate signal in (A), high signal.	91

List of Figures *(Cont...)*

Fig. No.	Title	Page No.
Fig. (59):	Axial T2 and Sagittal T2 of the left hand showing high signal showing a well-defined binoculars subcutaneous cystic swelling is observed opposite to posterior lateral aspect of distal phalanx of index finger.....	92
Fig. (60):	Coronal T1 (A), coronal T2 (B), coronal PDWI (D) and coronal STIR (C) of the left hand.....	93
Fig. (61):	Axial T1 (A), axial T2 (B) Sagittal T1 (C) and Sagittal T2 (D) of the right hand showing intermediate T2 and high T1 it is centered in the 2nd inter digital web spaces between the index and middle fingers.....	96
Fig. (62):	Coronal T1 (A), Coronal T2 (B) and Coronal STIR (C) showing intermediate signal in (A), high signal (B& C) it is centered in the 2nd inter digital web spaces between the index and middle fingers, expanding the deep palmar space, incasing the 2nd, third finger tendon, neurovascular bundles up to the wrist.....	97
Fig. (63):	Axial T1 (A), axial T2 (B) and Sagittal T2 (C) Soft tissue mass lesion is seen at the radial aspect of the right wrist joint of showing intermediate T2 and high signal T1, this lesion shows scalloping of the lateral aspect of the trapezium bone with small tongue projection seen between the bases of the 1st and 2nd metacarpal.....	99

List of Figures *(Cont...)*

Fig. No.	Title	Page No.
Fig. (64):	Coronal T1 (A), Coronal T2 (B) and Coronal PDWI(C) showing Soft tissue mass lesion is seen at the radial aspect of the right wrist joint of showing intermediate T2 and high signal T1 and PDWI, this lesion shows scalloping of the lateral aspect of the trapezium bone with small tongue projection seen between the bases of the 1st and 2nd metacarpal.....	100
Fig. (65):	Axial T1 (A), axial T2 (B) axial DWI (C), axial ADC (D) and Sagittal T2 (F) of right hand Expansile well defined corticated lesion seen involving the base of the proximal phalanx of the 5th finger eliciting iso intense to low signal on T1 and ADC bright signal. On T2 and DW it shows fine internal septation.it is seen involving the proximal meta-diaphyseal aspect of bone.....	102
Fig. (66):	Coronal T1 (A), Coronal T2 (B) and Coronal STIR (C) showing right hand Expansile well defined corticated lesion seen involving the base of the proximal phalanx of the 5th finger eliciting iso intense to low signal on T1, bright signal on T2 and STIR. It show fine internal septation involving the proximal meta-diaphyseal aspect of bone.....	103

List of Figures (Cont...)

Fig. No.	Title	Page No.
Fig. (67):	Axial T1 (A) and axial T2 (B) left index finger show a tiny lesion at subungual position of index finger, it seen at the medial aspect of the nail bed measuring 0.3x0.5 cm, bright signal in T2 and low signal in T1WSI.	105
Fig. (68):	Coronal T1 (A), Coronal T2 (B) and Coronal STIR (C) left index finger show a tiny lesion at subungual position of index finger, it seen at the medial aspect of the nail bed measuring 0.3x0.5 cm, bright signal in T2, STIR, low signal in T1WSI.	106
Fig. (69):	Axial T1 (A), axial T2 (B) axial STIR (C) and Sagittal T2 (D), The right middle finger shows a lobulated soft tissue seen between the proximal and middle phalanges and extending to the lateral aspect of the finger, seen eliciting low T1, T2 and high signal STIR. It shows deep extension beneath the flexor tendons of the middle finger.....	108
Fig. (70):	Coronal T1 (A), Coronal T2 (B) and Coronal STIR (C) The right middle finger shows a lobulated soft tissue seen between the proximal and middle phalanges and extending to the lateral aspect of the finger, seen eliciting high STIR signal and low signal on T1 and T2. It shows deep extension beneath the flexor tendons of the middle finger.	109